



rivers and
water quality

arteries of the Australian environment

LandWater & Wool
Shaping the future



another australian wool innovation limited

Rapid Appraisal of Riparian Condition

Technical guideline for the southern tablelands of New South Wales

Amy Jansen, Alistar Robertson, Leigh Thompson, Andrea Wilson and Fleur Flanery

Summary

- ~ Riparian habitats are where land and water ecosystems meet. They are vital sites in a catchment supporting high levels of biodiversity. This is true even for riparian areas adjacent to creeks and gullies that may flow with water only occasionally.
- ~ Given the extensive degradation of riparian zones in Australia, there is a need for a rapid method of measuring riparian condition to underpin strategies for improved management.
- ~ Riparian condition refers to the degree to which human-altered ecosystems diverge from local semi-natural ecosystems in their ability to support a community of organisms and perform ecological functions.
- ~ The Rapid Appraisal of Riparian Condition assesses the ecological condition of riparian habitats using indicators that reflect functional aspects of the physical, community and landscape features of the riparian zone. The index is made up of five sub-indices, each with a number of indicators: Habitat continuity and extent (HABITAT), Vegetation cover and structural complexity (COVER), Dominance of natives *versus* exotics (NATIVES), Standing dead trees, hollows, fallen logs and leaf litter (DEBRIS), and Indicative features (FEATURES).
- ~ The Rapid Appraisal of Riparian Condition has been used in south-eastern Australia to examine relationships between grazing intensity and riparian condition. Generally, poor riparian condition was associated with high levels of grazing intensity.
- ~ Testing of the Rapid Appraisal of Riparian Condition index confirms that it is a good indicator of the biodiversity and functioning of riparian zones.
- ~ The Rapid Appraisal of Riparian Condition has been trialled on ephemeral and permanent creek systems around Bookham and Yass on the southern tablelands of New South Wales.



Land, Water & Wool
is a joint Australian
Wool Innovation Ltd
and Land & Water
Australia program.

Background

Riparian habitats are where land and water ecosystems meet. They are vital places in a farm and in a catchment, supporting high levels of biodiversity and being critical in controlling flows of energy and nutrients between the land and the water (creeks and rivers) (Naiman & Decamps, 1997). This is true even on creeks and gullies that only occasionally carry flowing water. Riparian areas are powerful indicators of catchment quality (e.g. Rapport et al., 1998). Human settlement has always been focused on rivers and is often a major determinant of riparian structure and function (e.g. Dynesius & Nilsson, 1994). One of the biggest impacts on riparian areas has been the introduction of domestic stock, with grazing being the major land use over 60% of Australia's land surface (Wilson, 1990). Stock concentrate around water sources, which means riparian and wetland habitats, as well as those around artificial watering points in pastoral regions, suffer greater impacts from domestic and feral grazing herds than dryland areas (Robertson, 1997; James et al., 1999). These impacts have led to extensive loss of ecological condition in riparian areas in Australia.

To improve the management of riparian areas and help land managers decide on the priorities for management of their streams or creeks, baseline assessments of the riparian condition and the factors that have contributed to it should be undertaken. An effective and simple method for doing this has been developed based on rapid appraisal techniques to measure ecosystem condition or integrity (Fairweather, 1999; Boulton, 1999).

The Rapid Appraisal of Riparian Condition index can provide a comparison of areas on-farm and with local areas considered to be in natural condition; it helps to pinpoint problems where management intervention is required. This method is suitable for periodic repeat assessments in the same area so that changes can be tracked over time and management adjusted if necessary (an adaptive management approach). This Guideline describes the Rapid Appraisal of Riparian Condition tailored for the southern tablelands region of New South Wales.

Throughout this Guideline, riparian condition refers to the degree to which human-altered ecosystems diverge from local semi-natural ecosystems in their ability to support a community of organisms and perform ecological functions (c.f. Karr, 1999).

Rapid Appraisal of Riparian Condition (RARC)

Assessment methods incorporating indicators of geophysical and biological properties and processes are likely to provide reliable estimates of ecological condition in riverine ecosystems (Fairweather, 1999; Boulton, 1999). Ladson et al. (1999) described an index of stream condition based on 18 indicators that measure alterations to the hydrology, physical form, streamside vegetation, water quality and biota of streams. This project used a similar approach, and chose indicators to reflect functional aspects of the physical, community and landscape features of the riparian zone, as defined by Naiman and Decamps (1997) (see Table 1). Some of the indicators chosen reflect the variety of functions, e.g. different aspects

of vegetation cover play a role in reducing bank erosion, providing organic matter and habitat for fauna, and providing connections in the landscape. The RARC index is made up of five sub-indices, each with a number of indicator variables (see Table 2, overleaf). The indices cover:

1. Habitat continuity and extent (HABITAT).
2. Vegetation cover and structural complexity (COVER).
3. Dominance of native *versus* exotic plants (NATIVES).
4. Standing dead trees, fallen logs and leaf litter (DEBRIS).
5. Indicative features (FEATURES).

Table 1. Summary table of functions, components and indicators assessed in the Rapid Appraisal of Riparian Condition index.

Functions of the riparian zone at different levels of organisation	Components of the riparian ecosystem that perform those functions	Indicators of the functions used in the RARC
<i>Physical:</i>		
Reduction of erosion of banks	Roots, ground cover	Vegetation cover*
Sediment trapping	Roots, fallen logs, ground cover	Canopy cover, fallen logs, ground cover vegetation, leaf litter** cover
Controlling stream microclimate/ discharge/water temperatures	Riparian forest	Canopy cover
Filtering of nutrients from upslope	Vegetation, leaf litter	Ground cover vegetation, leaf litter cover
<i>Community:</i>		
Provision of organic matter to aquatic food chains	Vegetation	Vegetation cover, leaf litter cover
Retention of seeds, bulbs, stems and other sources of natural plant regeneration	Fallen logs, leaf litter	Fallen logs, leaf litter cover
Maintenance of plant diversity	Regeneration of dominant species, presence of important species, dominance of natives <i>versus</i> exotics	Native canopy and shrub regeneration, grazing damage to regeneration, reeds, native vegetation cover
Provision of habitat for aquatic and terrestrial fauna	Fallen logs, leaf litter, standing dead trees/hollows, riparian forest, habitat complexity	Fallen logs, leaf litter cover, standing dead trees, vegetation cover, number of vegetation layers
<i>Landscape:</i>		
Provision of biological connections in the landscape	Riparian forest (cover, width, connectedness)	Vegetation cover, width of riparian vegetation, longitudinal continuity of riparian vegetation
Provision of refuge in droughts	Riparian forest	Vegetation cover

* Vegetation cover = canopy, understorey and ground cover.

** Leaf litter includes any dead plant material such as leaves, grasses, twigs and bark.

Table 2. Sub-indices and indicators of the Rapid Appraisal of Riparian Condition, the range within which each is scored, the method of scoring for each indicator, and the maximum possible total for each sub-index.

Sub-index	Indicator	Range	Method of scoring	Total
HABITAT				11
	Longitudinal continuity of riparian vegetation (≥ 5 m wide)	0–4	0 = $< 50\%$, 1 = 50–64%, 2 = 65–79%, 3 = 80–94%, 4 = $\geq 95\%$ vegetated bank; with 1/2 point subtracted for each significant discontinuity (≥ 50 m long)	
	Width of riparian vegetation (scored differently for channels $<$ or ≥ 10 m wide)	0–4	<i>Channel ≤ 10 m wide:</i> 0 = VW < 5 m, 1 = VW 5–9 m, 2 = VW 10–29 m, 3 = VW 30–39 m, 4 = VW ≥ 40 m <i>Channel > 10 m wide:</i> 0 = VW/CW < 0.5 , 1 = VW/CW 0.5–0.9, 2 = VW/CW 1–1.9, 3 = VW/CW 2–3.9, 4 = VW/CW ≥ 4 , where CW = channel width and VW = vegetation width	
	Proximity to nearest patch of intact native vegetation > 10 ha	0–3	0 = > 1 km, 1 = 200 m–1 km, 2 = contiguous, 3 = contiguous with patch > 50 ha	
COVER				12
	Canopy (> 5 m tall)	0–3	0 = absent, 1 = 1–30%, 2 = 31–60%, 3 = $> 60\%$ cover	
	Understorey (1–5 m tall)	0–3	0 = absent, 1 = 1–5%, 2 = 6–30%, 3 = $> 30\%$ cover	
	Ground (< 1 m tall)	0–3	0 = absent, 1 = 1–30%, 2 = 31–60%, 3 = $> 60\%$ cover	
	Number of layers	0–3	0 = no vegetation layers to 3 = ground cover, understorey and canopy layers	
NATIVES				9
	Canopy (> 5 m tall)	0–3	0 = none, 1 = 1–30%, 2 = 31–60%, 3 = $> 60\%$ cover of native plants	
	Understorey (1–5 m tall)	0–3	0 = absent, 1 = 1–5%, 2 = 6–30%, 3 = $> 30\%$ cover of native plants	
	Ground (< 1 m tall)	0–3	0 = none, 1 = 1–30%, 2 = 31–60%, 3 = $> 60\%$ cover of native plants	
DEBRIS				10
	Leaf litter	0–3	0 = none, 1 = 1–30%, 2 = 31–60%, 3 = $> 60\%$ ground cover	
	Native leaf litter	0–3	0 = none, 1 = 1–30%, 2 = 31–60%, 3 = $> 60\%$ ground cover	
	Standing dead trees (> 20 cm dbh)	0–1	0 = absent, 1 = present	
	Hollow-bearing trees	0–1	0 = absent, 1 = present	
	Fallen logs (> 10 cm diameter)	0–2	0 = none, 1 = small quantities, 2 = abundant	

dbh = diameter at breast height, $<$ less than, \leq less than or equal to, $>$ greater than, \geq greater than or equal to.

Table 2. continued

Sub-index	Indicator	Range	Method of scoring	Total
FEATURES				8
	Native canopy species regeneration (< 1 m tall)	0–2	0 = none, 1 = scattered, 2 = abundant; with 1/2 point subtracted for grazing damage	
	Native understorey regeneration	0–2	0 = none, 1 = scattered, 2 = abundant; with 1/2 point subtracted for grazing damage	
	Large native tussock grasses	0–2	0 = none, 1 = scattered, 2 = abundant	
	Reeds	0–2	0 = none, 1 = scattered, 2 = abundant	



Photo 1. A site in excellent condition on the Upper Murrumbidgee River (RARC score = 46; this site includes continuous vegetation along the riparian zone, standing dead trees and fallen logs, native shrub understorey and regeneration of canopy trees). Photo Greening Australia.



Photo 2. A site in very poor condition on the Murrumbidgee River from Tamar's Bridge (RARC score = 12; note limited native overstorey, lack of understorey and no native regeneration). Photo Greening Australia.

Photos 1 and 2 show contrasting sites in excellent and very poor condition. Details of the scoring for these sites can be found in the box below.

Example of scoring indicators for the sites shown in Photos 1 and 2 (see Table 2 for indicators and details)				
Sub-index	Excellent condition site (Photo 1)		Very poor condition site (Photo 2)	
Habitat	4 + 4 + 3 =	11	0 + 0 + 0 =	0
Cover	3 + 3 + 3 + 3 =	12	1 + 0 + 3 + 2 =	6
Natives	3 + 3 + 3 =	9	1 + 0 + 1 =	2
Debris	3 + 3 + 1 + 0 + 1 =	8	1 + 1 + 1 + 0 + 0 =	3
Features	2 + 1 + 2 + 1 =	6	1 + 0 + 0 + 0 =	1
Total		46		12

Application of the Rapid Appraisal of Riparian Condition index

The RARC was initially developed as a tool to determine the impacts of grazing management practices on riparian condition, and to identify those practices which resulted in minimal impacts. We have now tested this approach in three areas of south-eastern Australia (see Figure 1); some results are presented below. Note that these results were obtained using the original version of the RARC rather than the version modified for southern NSW, but the two versions give very similar scores.






Figure 1. Location of sites where the RARC has been applied.


Murrumbidgee River

A total of 138 sites (each 1 kilometre in length) were surveyed between Gundagai and Hay, on private properties, crown land and State Forests (Jansen & Robertson, 2001a). The majority of sites on private property were in very poor condition, while sites on Crown Land (mainly Travelling Stock Reserves) were very variable. Most State Forest sites were in good to excellent condition (Figure 2a).



Gippsland

A total of 108 sites (each 150 metres in length) were surveyed in West and South Gippsland, at three types of sites — grazed paddocks on private properties, planted and fenced riparian areas on private properties, and remnant patches of uncleared native vegetation both on private properties and in reserves (Thompson et al., 2003). All private property sites were on dairy farms. The majority of sites were in very poor condition, with only remnant sites scoring above average (Figure 2b). It should be noted that most planted sites were relatively recently fenced, and their condition can be expected to improve as the plantings mature.



Goulburn-Broken

A total of 46 sites (each 200 metres in length) were surveyed in the upper and mid-Goulburn-Broken catchment, at grazed and ungrazed sites on private properties, and at ungrazed sites in reserves (Wilson et al., 2003). Again, the majority of sites were in very poor condition (Figure 2c). Like the Gippsland planted sites, many of the Goulburn-Broken ungrazed sites on private properties were relatively recently fenced, and their condition can be expected to improve as plantings mature.




Figure 2. The number of sites scoring in each category (< 25 very poor, 25–30 poor, 30–35 average, 35–40 good and > 40 excellent) of the RARC index for three regions: (a) Murrumbidgee River, (b) West and South Gippsland, and (c) upper and mid-Goulburn-Broken catchment.

Riparian condition in relation to stocking rates

In all three regions, we examined the relationship between stocking rates and riparian condition, with Figure 3 below showing our results. Clearly, riparian condition declined with increased stocking rates, across all regions and a large range of stocking rates. Given the large number of sites in poor condition in all catchments, this suggests that stocking rates commonly used on private properties are too high to maintain riparian zones in good condition.




Figure 3. RARC condition scores in relation to stocking rates (DSE/ha/annum) for three regions: Murrumbidgee River, West and South Gippsland, and upper and mid-Goulburn-Broken catchment.

RARC assessments in the southern tablelands

A total of 26 sites around Yass and Bookham were surveyed using the Rapid Appraisal of Riparian Condition tool (RARC) between March 2005 and March 2006 by the Land, Water & Wool project officer Fleur Flanery. (See other useful publications listed on pages 14 and 15.)

The majority of RARC assessments were conducted on smaller creeks on farms in the Yass River sub-catchment of the Murrumbidgee catchment between March and June 2005. One assessment was conducted at the Yass Weir which is being revegetated and currently has little native vegetation and another was conducted on Crown Land near the township's main weir. All surveys were taken along a 200 metre section of the waterway. The scores obtained at a selection of these sites are shown in the table below.

The majority of the sites assessed using the RARC were in poor condition with scores ranging from 8 to 28. Many sites had some tree cover and ground cover, but the understorey of native shrubs, grasses and forbs was absent from all sites. Several sites had little or no native riparian vegetation. Sites scoring 30 or less are generally in poor to very-poor ecological condition. The majority of sites on private land showed riparian areas in poor condition resulting from continual grazing which reduced the understorey along the banks and inhibited natural regeneration.

Table 3. RARC scores from a range of sites surveyed within the Yass–Bookham region between March 2005 and March 2006.

Site	RARC score (max. 50)	Comments about site condition
Bango Creek	17.5	Poor condition. Some existing riparian vegetation but no natural regeneration.
Talmo Creek	21.0	Poor condition. Native riparian vegetation is present but no regeneration visible.
Yass River Weir	10.4	Very poor condition. Limited vegetation of any type and no sign of regeneration.
Eight Mile Creek	28.0	Average condition. Recreational area, ungrazed, but mown, some natural regeneration visible.
Chinamen's Creek	14.7	Poor condition. Dominated by exotic species.
Gums Lane Creek	28.0	Average condition. Dominated by native species with natural regeneration visible.
Back Creek, Yass	7.9	Very poor condition. Little or no riparian vegetation.
Cattleyard Creek, Harden	11.8	Poor condition. Good ground cover but no trees or natural regeneration visible.

This is one of the summary tables prepared by the project officer following RARC assessments of each site. The table would be more informative if details were provided about the type of land use and grazing regime being used in each of the sites, as management actions can then be linked to riparian health. This information can also provide the basis upon which new management strategies are developed to improve riparian condition.

Sub-indices of the riparian condition index

There was variation across regions in relation to which sub-indices accounted for most of the variation in the total riparian condition score (Table 4). In the Murrumbidgee region, 85% of the variance in the total condition score was explained by the DEBRIS sub-index (scoring for leaf litter, fallen logs and standing dead trees). In Gippsland, 90% of the variance in the total condition score was explained by the NATIVES sub-index (scoring for native species in the vegetation cover and debris). In the Goulburn-Broken, 79% of the variance in the total condition score was explained by the COVER sub-index (scoring for % cover in each vegetation layer, and the number of vegetation layers).

The DEBRIS sub-index consistently explained at least 70% of the variance in the total condition score, suggesting that management practices aimed at retaining standing dead trees and fallen logs would improve riparian condition scores in all regions. The HABITAT sub-index was also relatively consistent across regions, explaining at least 62% of the variance in total condition scores. This suggests that maintaining or restoring a continuous canopy in the riparian zone is also important in all regions. In contrast, the NATIVES sub-index explained little of the variance in the Murrumbidgee but most of it in Gippsland. This sub-index indicates that in the Murrumbidgee, the canopy trees are predominantly native, there is little shrub cover, and the ground cover is predominantly exotic. In this region, there is little chance of altering this on a large scale. In Gippsland, however, the index indicates a lot of variability in the dominance of natives over exotics in all vegetation layers, and that management aimed at maintaining or restoring native species could significantly improve riparian condition.

Sub-index	Murrumbidgee	Gippsland	Goulburn-Broken
COVER	0.42	0.83	0.79
DEBRIS	0.85	0.75	0.70
HABITAT	0.81	0.80	0.62
NATIVES	0.23	0.90	0.77
FEATURES	0.60	0.32	0.56

Table 4. Proportion of variance in the total riparian condition index score explained (R^2 value) by each sub-index for three regions: Murrumbidgee River, West and South Gippsland, and upper and mid-Goulburn-Broken catchment. The R^2 value was obtained by regressing the values for each sub-index against the total index scores for each site.

Why is the RARC a useful tool?

What does riparian condition tell us about the biodiversity and functioning of riparian zones?

The RARC has been tested against more detailed measures of the biodiversity and functioning of riparian zones in the Murrumbidgee and Gippsland regions. There was a significant positive relationship between litter decomposition rates in the soil and the COVER sub-index of the RARC score in both Summer ($r = 0.50$, $p < 0.05$) and Autumn ($r = 0.78$, $p < 0.01$), indicating that decomposition rates were higher where there was more vegetation cover in the riparian zone of the Murrumbidgee River (Robertson, Wassens & Jansen, in prep.). There were highly significant relationships between bird communities and all sub-indices, as well as the total RARC score ($r = 0.68$, $p < 0.0001$), indicating that riparian bird communities varied according to the condition of the riparian zone of the Murrumbidgee River (Jansen & Robertson, 2001b). Of particular significance ($r = 0.74$, $p < 0.0001$) was the DEBRIS sub-index (scoring for leaf litter, fallen logs and standing dead trees), indicating that retention of leaf litter and woody debris in riparian habitats is crucial to the survival of riparian bird communities. Many of the species most dependent on these features (e.g. Brown Treecreepers) are threatened or declining throughout the agricultural regions of southern Australia (Ford et al., 2001).

In Gippsland, there was also a significant relationship ($r = 0.59$, $p < 0.0001$) between bird communities and the total RARC score, indicating again that riparian bird communities varied according to the condition of riparian zones in Gippsland (Thompson et al., 2003).

Given the importance of riparian zones in supporting high levels of regional biodiversity (Naiman & Decamps, 1997), and the links between riparian condition and biodiversity demonstrated here, the RARC is a useful tool for assessing riparian condition and hence biodiversity and functioning of riparian zones.

r = correlation coefficient
(indicates the strength of a relationship)

p = significance
(where $p < 0.05$ indicates a significant relationship)



This creek is in excellent condition with an intact understorey, logs and other debris for wildlife habitat and shade over the water. Stock are not allowed into the area. Photo Lori Gould.



The Clamorous Reed Warbler. These birds live in riparian areas and their presence may be used as an indicator of riparian health. Photo Julian Robinson.

Applying the RARC: Steps in assessing riparian condition

The Rapid Appraisal of Riparian Condition index can be used for a variety of applications. Examples include determining relationships between riparian condition and management practices, as in the studies described earlier, or surveying overall condition within a catchment to determine priorities for future rehabilitation works in the catchment. Whatever the application, care should be taken to clearly define the question to be answered, determine the sampling design and select sites appropriately to answer the question. This may require help from a consultant with experience in experimental design and data analysis. In general, sampling of sites should be *random**, rather than only sampling sites which are easily accessible by road.

A single observer should conduct all assessments, and they should undertake some training beforehand, to ensure consistency of data collection. The observer will need to have some experience in discriminating native and exotic plant species, and may benefit from previous experience in habitat surveys.

All sites should be surveyed at a similar time of year. Use a separate scoring sheet for each site. Allow 20–60 minutes per site, depending on size and accessibility.

*If you were interested in surveying overall catchment condition, you could choose sites randomly by laying a grid over a map of the catchment, locating and numbering all squares which contain a riparian zone, then putting these numbers in a hat and pulling out as many sites as you wish to sample.

1 Determine site size

Site size must be determined according to the size of the management unit of interest. For example, our studies have examined impacts of grazing management on riparian condition, so management units have been individual paddocks. On the Murrumbidgee River, where paddocks are relatively large, a 1 kilometre length of the riparian zone was defined as a 'site', while in Gippsland, where paddocks are much smaller, a 150 metre length was used. Ideally, sites should be at least 200 metres long, with 500 metres being the preferred length where practicable. On larger rivers, only one side of the river is surveyed, while at smaller sites where it is practicable to do so, both sides may be surveyed (provided they are subject to the same management regime).



Two sections of the Yass River approximately 2 kilometres apart. Riparian areas can change dramatically depending on their management. To get an accurate assessment of the general condition of the riparian zone along a river, several RARC assessments should be conducted at randomly selected sites. Both these riparian areas are in poor condition due to stock grazing and the vegetation being cleared from along the banks. The RARC assessment can assist woolgrowers work out which management strategies can be used in these situations to improve riparian and in-stream health. Photos Greening Australia.

The transects at each site should ideally traverse the width of the riparian zone. However, this is not always easy to determine in the field. To simplify this, we use a transect length determined by the width of the river channel — 40 metres long for channels < 10 metres wide, and four times the channel width for larger rivers. A minimum width of 40 metres should be assessed, unless there is a very clear distinction between riparian and non-riparian areas. Where the riparian zone is clearly narrower than 40 metres or four times the channel width (for example, in a gorge), the transect length should be adjusted accordingly. Where the riparian zone is much wider than this (for example, on a lowland floodplain river), four times the channel width should be adequate to represent the riparian zone. Figure 4 illustrates a hypothetical river with the layout of the survey area and the transects indicated.

2 Score indicators

A sample scoring sheet can be found on page 16 of this Guideline. The complete scoring system is summarised in Table 2. Longitudinal continuity and proximity are given single values for the whole site. All other indicators are scored along four transects (10 metres wide; perpendicular to the direction of river flow) evenly spaced along the bank.

HABITAT

Longitudinal continuity of riparian canopy vegetation: At each site, canopy vegetation along the bank is mapped to show the length and number of any discontinuities (gaps of more than 50 metres) in canopy cover (the bank is considered to be vegetated if the riparian canopy vegetation is at least 5 metres wide). Longitudinal continuity is then scored as follows:

0 = < 50%, 1 = 50–64%, 2 = 65–79%, 3 = 80–94%,
4 = ≥ 95% vegetated bank; with 1/2 point subtracted
for each significant discontinuity (> 50 m long)

Proximity: An assessment is made of the shortest distance to the nearest patch of at least 10 hectares of relatively intact native vegetation (with an extra point if the area being assessed is within a patch of at least 50 hectares of relatively intact native vegetation). This can be assessed on-site or later using aerial photographs. Proximity is then scored as follows:

0 = > 1 km, 1 = 200 m–1 km, 2 = contiguous,
3 = contiguous with patch > 50 ha




Figure 4. Hypothetical river with length and transects marked. The scoring for the indicators in this diagram is shown (see page 16 for full score sheet).

A patch of relatively intact native vegetation should have at least the dominant overstorey vegetation remaining. This may not be trees, if the area is a natural grassland or shrubland.

For example, for a channel 12 metres wide and a vegetation width of 30 metres, $VW/CW = 2.5$, giving a score of 3.

Width of riparian vegetation: The channel width is defined by the area within the banks that is normally lacking any terrestrial or bankside vegetation. The width of the riparian canopy vegetation is the distance from the bank to the first gap of > 50 metres in the canopy vegetation. Channel width (CW) and width of the riparian vegetation (VW) are estimated to the nearest 5 metres in the field. For channels less than 10 metres wide, the vegetation width is converted directly to a score, while for channels more than 10 metres wide, the vegetation width is divided by the channel width to obtain the score as follows:

Channel ≤ 10 m wide: 0 = $VW < 5$ m, 1 = $VW 5-9$ m, 2 = $VW 10-19$ m, 3 = $VW 20-39$ m, 4 = $VW \geq 40$ m

Channel > 10 m wide: 0 = $VW/CW < 0.5$, 1 = $VW/CW 0.5-0.9$, 2 = $VW/CW 1-1.9$, 3 = $VW/CW 2-3.9$, 4 = $VW/CW \geq 4$

COVER (see photo 3 overleaf)

Vegetation cover within each layer is scored as follows:

Ground cover (lichens, mosses, grasses, herbs, reeds and sedges to 1 m tall): 0 = none, 1 = 1-30%, 2 = 31-60%, 3 = > 60%

Understorey cover (herbs, reeds, shrubs and saplings 1-5 m tall): 0 = none, 1 = 1-5%, 2 = 6-30%, 3 = > 30% (Note that understorey cover is scored on a different scale to the others, since it is normally less dense.)

Canopy cover (trees > 5 m tall): 0 = none, 1 = 1-30%, 2 = 31-60%, 3 = > 60%

The number of layers of vegetation is scored as follows:

0 = no vegetation layers to 3 = ground cover, understorey and canopy layers



Photo 3. Canopy cover increasing from 1 to 3 (left to right). Photos Amy Jansen.



Tussocky perennial (long-lived) grasses tend to be native species while annual (short-lived) grasses tend to be exotic species (with a few obvious exceptions such as *Phalaris* which is a perennial exotic species).

Photo 4. Exotic annual ground cover (left) versus native perennial tussock ground cover (right). Photos Amy Jansen.



Photo 5. Leaf litter cover increasing from 1 to 3 (left to right). Photos Amy Jansen.

NATIVES (see photo 4 above)

Native vegetation cover within each layer is scored as for cover, but excluding the contribution of exotic species (to estimate cover of native species, imagine removing all exotic species and re-estimating vegetation cover with only the native species):

Ground cover (lichens, mosses, grasses, herbs, reeds and sedges to 1 m tall): 0 = none, 1 = 1–30%, 2 = 31–60%, 3 = > 60%

Understorey cover (herbs, reeds, shrubs and saplings 1–5 m tall): 0 = none, 1 = 1–5%, 2 = 6–30%, 3 = > 30%

Canopy cover (trees > 5 m tall): 0 = none, 1 = 1–30%, 2 = 31–60%, 3 = > 60%

DEBRIS (see photo 5 above)

Cover of leaf litter on the ground, and cover of native leaf litter are scored as follows:

0 = none, 1 = 1–30%, 2 = 31–60%, 3 = > 60% cover

Standing dead trees > 20 cm diameter at breast height, and hollow-bearing trees (look for dead branches and broken-off branch stubs in large trees which may have developed hollows) are scored as follows:

0 = absent, 1 = present

Fallen logs (>10cm diameter) are scored as follows:

0 = none, 1 = small quantities, 2 = abundant (where small quantities = one or two logs, and abundant = three or more logs)

FEATURES

The abundances of native canopy species regeneration (< 1 metre tall) and native understorey regeneration are scored as follows:

0 = none, 1 = scattered, and 2 = abundant, with 1/2 point subtracted for grazing damage (where scattered = one or two seedlings, and abundant = three or more seedlings; grazing damage is evidence that any of the seedlings have been browsed by grazing animals such as domestic livestock or kangaroos)



Photo 6. *Poa labillardieri*, an example of a large native tussock grass found in riparian zones. Photo Amy Jansen.

The abundances of large native tussock grasses (species such as *Poa labillardieri*) and reeds (species such as *Phragmites*, *Typha* (Cumbungi) and *Carex* which are normally only found on riverbanks or in swampy areas) are scored as follows:

0 = none, 1 = scattered, and 2 = abundant (where scattered = one or two plants, and abundant = three or more plants)

3 Analyse data

The indicators are averaged across transects, then summed into sub-indices. The final index score is then the sum of the sub-indices, with a possible maximum of 50 indicating best condition. To examine the results, it is helpful to categorise the index scores, e.g. less than 25 very poor, 25–30 poor, 30–35 average, 35–40 good and more than 40 excellent. It is also helpful to examine sub-index scores, and to determine which sub-indices contribute most to the final condition score. This can be done by regression of sub-index scores on the total index score.

4 Benchmarking

The scoring system given here has been developed for a generalised riparian area in south-eastern Australia, and may need to be adjusted for particular situations. Ideally, a number of relatively pristine sites in the region should be surveyed to provide a benchmark for the scoring system. The scores for each indicator can then be checked to ensure that all indicators are present, and that the maximum score can be achieved for each indicator. For example, in wet forests with a dense canopy, there may be no large tussock grasses but ferns could be used as an indicator instead. Also, ground cover may never reach > 60% due to shading, so this indicator may need to be adjusted accordingly (for example, the scores given for different levels of ground cover could be rescaled similarly to those given for understorey cover). Benchmarking against relatively pristine sites is not always possible in highly modified catchments. In these situations, we can only make a ‘best guess’, based on local knowledge and historical information, about the appropriate scoring for each indicator in these catchments.

5 Limitations of the RARC

While the RARC outlined in this booklet has been tested in a number of catchments and situations, it has some limitations:

- ~ The RARC that is presented in this publication has been designed and tested on creeks and rivers in south-eastern Australia that are naturally dominated by trees, with at least 60% canopy cover. Three modified versions of the RARC have been developed to suit particular regions. The version for tropical savannas is called the Tropical Rapid Assessment of Riparian Condition (TRARC). The mid north of South Australia has a RARC that has been developed for riparian areas that don’t have overhead trees and people in the Tasmanian midlands can refer to a RARC that accounts for particular vegetation types found along their riparian zones. For copies of these RARCs and the scoring sheets that accompany them go to www.rivers.gov.au
- ~ The RARC is intended as an indicator of *current* condition. This means that for restored areas, it will not indicate the potential for recovery of ecosystem function.

Further information

We will be continuing to refine and update the RARC, so to get the most recent version check — www.rivers.gov.au/lww. There is an Excel spreadsheet available on the website which includes a copy of a field data sheet for printing, and a data entry sheet. If you enter the data for a site, it will automatically calculate the averages for each transect and the final sub-index and total scores for you. If you have a number of sites, you will need to save a separate copy of the worksheet for each site. There is also a field calculation sheet which you can print on the reverse of the field data sheet if you wish to calculate scores as you go in the field (you may need a calculator to take the averages across the transects).

The method was developed by Amy Jansen, Alistar Robertson, Leigh Thompson and Andrea Wilson. It is also available as *River and Riparian Technical Update*, no. 4A from Land & Water Australia. Go to www.rivers.gov.au for details.

Good reads for primary producers who want to know more about their waterways



Managing in-stream wetlands on wool-producing farms — this fact sheet provides information about the benefits of, and management tips for, in-stream wetlands.
Product code PF061168



Managing gullies on wool-producing farms — this fact sheet provides information about gullies, their prevention and management, so that woolgrowers can make informed decisions about what to do on their farm.
Product code PF061168

Managing creeks and waterways in the southern tablelands of New South Wales: A woolgrowers guide — this is a summary of key management recommendations particular to the southern tablelands region. Packed with information, it is a "must have" for tablelands woolgrowers.
Product code PX071293



People who can help

Land, Water & Wool — NSW Rivers Project

Fleur Flanery

Project Officer (RARC trained)
Tel: 02 6281 8585 / 0408 627 774
E-mail: fflanery@greeningaustralia.org.au

Murrumbidgee Catchment Management Authority

Brad Parker

Catchment Project Officer (RARC trained)
Tel: 02 6118 6060
E-mail: brad.parker@cma.gov.au

Kathleen Harvey

Catchment Project Officer (RARC trained)
Tel: 02 6118 6060
E-mail: kathleen.harvey@cma.gov.au

Louise Hufton

Community Support Officer
Harden-Murrumburrah Landcare Group
Tel: 02 6386 3954
E-mail: louise.hufton@cma.gov.au

Greening Australia — Capital Region

Lori Gould (RARC trained)

Tel: 02 6253 3035
E-mail: lgould@act.greeningaustralia.org.au

NSW Department of Primary Industries

Phil Graham

Tel: 02 6226 2199
E-mail: phillip.graham@agric.nsw.gov.au

Organisations

Land & Water Australia

Tel: 02 6263 6000
Web: www.lwa.gov.au

Greening Australia

Tel: 1300 886 589
E-mail: exchange@greeningaustralia.org.au
Web: www.greeningaustralia.org.au

Publications for woolgrowers

The *Wool industry river management guides* bring together the latest science and recommended management practices for riparian areas within the context of a commercial wool growing property. The Guides are available for the high rainfall regions (above 600 mm) and sheep/wheat regions (300–600 mm) of Australia. Each book has over 200 full-colour pages.

In addition www.rivers.gov.au/lww will offer an active contents list which will give you a snapshot of what is in each section.

High rainfall zone: product code PX050951

Sheep/wheat zone: product code PX050952



Managing rivers, streams and creeks: A woolgrowers guide — is a summary of the key recommendations from the ‘Wool industry river management guides’ and provides an introduction to river and riparian management issues on farm.

Product code PX051003



Are my waterways in good condition? — a checklist that provides colour coded pictures that you can use to assess the condition of your stream or creek. It is a quick and easy way to work out the health of the streams or creeks running through your property, and it suggests management actions to improve or maintain these vital parts of your farm.

Product code PB061114




River insights — a publication featuring the stories of ten woolgrowers and what has motivated them to manage their rivers, creeks and streams in ways that make both economic and environmental sense.

Product code PK050950

Stock and waterways: a manager's guide — offers practical advice on how stock farmers can manage riparian land both productively and sustainably, and includes a number of case studies from farmers throughout Australia who have seen the benefits of changing their management practices.

Product code PR061132



These products are available from CanPrint Communications on freecall 1800 776 616 in hard copy, or can be downloaded from — www.landwaterwool.gov.au or www.rivers.gov.au

Rapid Appraisal of Riparian Condition

Site: _____ Site number: _____ GPS start: _____

Date: _____ Observer: _____ GPS end: _____

Longitudinal continuity of riparian canopy vegetation (> 5 m wide)

Map	Score

0 = < 50%, 1 = 50–64%, 2 = 65–79%, 3 = 80–94%, 4 = > 95% vegetated bank; with 1/2 point subtracted for each significant discontinuity (> 50 m long)

Width of riparian canopy vegetation

Transect	Channel Width (CW)	Vegetation Width (VW)	Score
1			
2			
3			
4			
Average			

Proximity

Score

Nearest patch of native vegetation > 10 ha:
 0 = > 1 km, 1 = 200 m–1 km,
 2 = contiguous, 3 = contiguous with patch > 50 ha

Channel < 10 m wide: 0 = VW < 5 m, 1 = VW 5–9 m, 2 = VW 10–19 m, 3 = VW 20–39 m, 4 = VW > 40 m vegetated

Channel > 10 m wide: 0 = VW/CW < 0.5, 1 = VW/CW 0.5–0.9, 2 = VW/CW 1–1.9, 3 = VW/CW 2–3.9, 4 = VW/CW > 4

Vegetation cover: Canopy > 5 m, Understorey 1–5 m, Ground cover < 1 m

Transect	Canopy	Native canopy	Understorey	Native understorey	Ground cover	Native ground cover	Number of layers
1							
2							
3							
4							
Average							

Canopy and ground cover: 0 = none, 1 = 1–30%, 2 = 31–60%, 3 = > 60%

Understorey cover: 0 = none, 1 = 1–5%, 2 = 6–30%, 3 = > 30%

Debris

Transect	Leaf litter	Native leaf litter	Standing dead trees	Hollow-bearing trees	Fallen logs
1					
2					
3					
4					
Average					

Leaf litter and native leaf litter cover: 0 = none, 1 = 1–30%, 2 = 31–60%, 3 = > 60%

Standing dead trees (> 20 cm dbh) and hollow-bearing trees: 0 = absent, 1 = present

Fallen logs (> 10 cm diameter): 0 = none, 1 = small quantities, 2 = abundant

Features

Transect	Native canopy species regeneration	Native understorey regeneration	Large native tussock grasses	Reeds
1				
2				
3				
4				
Average				

Regeneration < 1 m tall: 0 = none, 1 = scattered, and 2 = abundant, with 1/2 point subtracted for grazing damage

Reeds and large tussock grasses: 0 = none, 1 = scattered, and 2 = abundant

Calculation of scores

Site number: _____

Longitudinal continuity of riparian canopy vegetation

Score	A
-------	---

Width of riparian canopy vegetation

Average	B
---------	---

Proximity

Score	C
-------	---

Vegetation cover

	Canopy	Native canopy	Understorey	Native understorey	Ground cover	Native ground cover	Number of layers
Average	D	H	E	I	F	J	G

Debris

	Leaf litter	Native leaf litter	Standing dead trees	Hollow-bearing trees	Fallen logs
Average	K	L	M	N	O

Features

	Native canopy species regeneration	Native understorey regeneration	Large native tussock grasses	Reeds
Average	P	Q	R	S

Totals

Site number	Habitat	Cover	Natives	Debris	Features	Total
(out of)	11	12	9	10	8	50
	A+B+C	D+E+F+G	H+I+J	K+L+M+N+O	P+Q+R+S	



References

- Boulton, A.J. 1999. An overview of river health assessment: philosophies, practice, problems and prognosis. *Freshwater Biology*, vol. 41, pp. 469–479.
- Dynesius, M. & Nilsson, C. 1994. Fragmentation and flow regulation of river systems in the northern third of the world. *Science*, vol. 266, pp. 753–762.
- Fairweather, P.G. 1999. State of environment indicators of ‘river health’: exploring the metaphor. *Freshwater Biology*, vol. 41, pp. 211–220.
- Ford, H.A., Barrett, G.W., Saunders, D.A. & Recher, H.F. 2001. Why have birds in the woodlands of Southern Australia declined? *Biological Conservation*, vol. 97, pp. 71–88.
- James, C.D., Landsberg, J. & Morton, S.R. 1999. Provision of watering points in the Australian arid zone: a review of effects on biota. *Journal of Arid Environments*, vol. 41, pp. 87–121.
- Jansen, A. & Robertson, A.I. 2001a. Relationships between livestock management and the ecological condition of riparian habitats along an Australian floodplain river. *Journal of Applied Ecology*, vol. 38, pp. 63–75.
- Jansen, A. & Robertson, A.I. 2001b. Riparian bird communities in relation to land management practices in floodplain woodlands of south-eastern Australia. *Biological Conservation*, vol. 100, pp. 173–185.
- Karr, J.R. 1999. Defining and measuring river health. *Freshwater Biology*, vol. 41, pp. 221–234.
- Ladson, A.R., White, L.J., Doolan, J.A., Finlayson, B.L., Hart, B.T., Lake, S. & Tilleard, J.W. 1999. Development and testing of an Index of Stream Condition for waterway management in Australia. *Freshwater Biology*, vol. 41, pp. 453–468.
- Naiman, R.J. & Decamps, H. 1997. The ecology of interfaces: Riparian zones. *Annual Review of Ecology and Systematics*, vol. 28, pp. 621–658.
- Rapport, D.J., Gaudet, C., Karr, J.R., Baron, J.S., Bohlen, C., Jackson, W., Jones, B., Naiman, R.J., Norton, B. & Pollock, M.M. 1998. Evaluating landscape health: integrating societal goals and biophysical process. *Journal of Environmental Management*, 53, pp. 1–15.
- Robertson, A.I. 1997. ‘Land-water linkages in floodplain river systems: the influence of domestic stock’, in *Frontiers in Ecology: Building the Links* (N. Klomp & I. Lunt, Eds), pp. 207–218. Oxford: Elsevier Scientific.
- Thompson, L., Robertson, A., Jansen, A. & Davies, P. 2003. ‘Identifying Best Management Practices for Riparian Habitats in Gippsland Dairy Regions: Riparian Condition and Relationships with Farm Management’, *Johnstone Centre Report no. 178*. Wagga Wagga, NSW: Johnstone Centre, Charles Sturt University.
- Wilson, A.D. 1990. The effects of grazing on Australian ecosystems. *Proceedings of the Ecological Society of Australia*, vol. 16, pp. 235–244.
- Wilson, A., Jansen, A., Curtis, A. & Robertson, A. 2003. ‘Understanding Landholder Management of Riparian Zones in the Goulburn Broken Catchment’, *Johnstone Centre Report no. 177*. Wagga Wagga, NSW: Johnstone Centre, Charles Sturt University.

Rapid Appraisal of Riparian Condition, Technical guideline for the southern tablelands of New South Wales. Amy Jansen, Alistar Robertson, Leigh Thompson, Andrea Wilson and Fleur Flanery.

Published by Land, Water & Wool
GPO Box 2182 Tel: 02 6263 6000 E-mail: Land&WaterAustralia@lwa.gov.au
Canberra ACT 2601 Fax: 02 6263 6099 Website: www.landwaterwool.gov.au
www.rivers.gov.au

May 2007

© Land & Water Australia

All rights reserved. No part of this publication may be reproduced, stored in any retrieval system or transmitted in any form or by any means electronic, mechanical, photocopying, recording or otherwise without the prior written permission of the publisher.

The information in this publication has been published by Land & Water Australia to assist public knowledge and discussion and to help improve the sustainable management of land, water and vegetation. Where technical information has been provided by or contributed by authors external to the Corporation, readers should contact the author(s) and conduct their own enquiries before making use of that information.

ISBN Print 1 921253 40 1 ISBN Electronic 1 921253 46 0
Product number PF071292



Publication data

Jansen, A., Robertson, A., Thompson, L., Wilson, A. & Flanery, F. 2007, 'Rapid Appraisal of Riparian Condition, Technical guideline for the southern tablelands of New South Wales', Land & Water Australia, Canberra.

Cover photo Greening Australia. Design by Angel Ink, Canberra.
Printed by Paragon Printer, Canberra.